




The Cost of Manual Charting

The following editorial allows readers to discover the costs associated with manual charting to help build the business case for implementing an automated electronic interface through a return-on-investment method. This paper was authored by Becky Clarke, BS, RN and TELCOR EVP of Point of Care and published in Point of Care Journal.

At TELCOR, we are committed to providing industry-best point of care software solutions. For more than 30 years, we've continuously invested, grown, and worked to become leaders in healthcare software solutions, tools, and metrics to drive business outcomes.

Connecting manual results

A POCT system would not be complete without the ability to capture results for manually-performed tests. TELCOR offers WebMRE®, an add-on module that fully integrates with TELCOR QML® point-of-care solution to capture manual test results. That means all results — whether from connected devices or manually entered — are available within QML and interfaced to the LIS/EMR for consistent and consolidated point of care result reporting.



ABOUT BECKY CLARKE

Becky Clarke, B.Sc., RN has been an executive vice president of TELCOR since being founded in 1995. Prior to TELCOR, she has 20 years of clinical and information systems experience. Ms. Clarke's extensive knowledge of information, computing, integration technologies, and excellent workflow analysis skills allow her to successfully deploy clinical applications for effective and efficient healthcare delivery. She has had leading roles in design, deployment, and support of commercially successful information systems for radiology, laboratory, pharmacy, nursing and ancillary order entry, medication charting, and many state-of-the-art system interfaces. Throughout her career, she has been instrumental in strategic planning, tactical deployment, and building organizations required to support the manufacturing process and on-going market requirements of clinical information systems.

ABOUT TELCOR

TELCOR provides software solutions for point-of-care data management with LIS/EMR integration and revenue cycle management software and service for outreach and reference laboratories. All trademarks, service marks, trade names, trade dress, product names and logos appearing on this document are the property of their respective owners. Copyright 2015-2026 by TELCOR Inc. All rights reserved.

REACH FOR MORE >>>

866-489-1207

 telcor@telcor.com

Copyright 2026 TELCOR Inc. All rights reserved.



EDITORIAL

The Cost of Manual Charting

Becky Clarke, BS, RN

Abstract: Charting point-of-care results is generally not performed by the laboratory point-of-care team; however, the oversight of this activity remains the responsibility of the point-of-care program. Although point-of-care management and interfacing are not new, there remain a significant number of hospitals not yet utilizing automated electronic interfaces for charting of results to tests performed outside the clinical laboratory (also known as point of care). This article is designed to help quantify the costs associated with manual charting and help build the case for implementing an automated electronic interface through a return-on-investment method.

.....

Charting point-of-care results is generally not performed by the laboratory point-of-care team; however, the oversight of this activity remains the responsibility of the point-of-care program. Being intimately involved with the LifeScan exit from the hospital point-of-care glucose testing market has given me awareness into the number of programs where glucose results are still not automatically electronically interfaced to the laboratory information system/electronic medical record (LIS/EMR), and thus one can assume there is a manual process for documentation of these and other point-of-care testing results.

In this article, I will help to quantify the cost of manual charting, provide components for creating a return on investment (ROI) by investing in automated interfaces, and review the benefits of an electronic interface to the LIS/EMR for the point-of-care program. Not just for bedside glucose testing, but for all point-of-care testing including those tests performed manually. Let's begin with the cost of manual charting.

If results are not electronically interfaced, one can assume they are either documented onto a paper chart or manually entered into the EMR's nursing notes or flow sheet. Because charting of glucose results is similar to other nurse charting activities, a very interesting study, "Enhancing Patient Safety Through Electronic Medical Record Documentation of Vital Signs¹," studied both errors made manually charting into the EMR and onto a paper chart. The study found the error rate for manual entry into the EMR to be less than 5% compared with the paper chart, where the error rate was 10%.

The types of errors were both errors of omission and transcription. For the sake of this article, we are going to assume bedside glucose results are being manually entered into the EMR's nursing notes or flow sheet simply because we know the benefits of clinician access to electronic results as opposed to the single paper chart. Have you ever taken the time to quantify how much time is spent manually charting these results every day? All you need to know is the number of patient tests being performed, using glucose as our example, and an estimate of the number of minutes used to chart results for these tests. Once you have determined these variables, you can plug them into the simple formula to calculate the amount of time spent performing this manual activity.

Results/y	Results/d	min/Result	min/d	h/d
100,000	2/4	2	548	9

Taking this one step further, you can estimate the cost of this activity by taking the hours per day from the formula above and use your facility's average hourly rate to determine the total dollars spent.

h/d	Hourly Rate	\$/d	\$/y
9	\$40	\$360	\$131,400

REACH FOR MORE >>>

866-489-1207

 telcor@telcor.com



In the example above, using \$40 as the average hourly burdened rate, it calculates to a cost of \$131,400 each year to manually chart 100,000 patient glucose results in the EMR, assuming all were charted and charted correctly. Automated electronic interfaces remove this labor component. Not only will they remove this labor component, but they also provide the following other benefits:

- Eliminate omissions and transcription errors discussed in the previously referenced article.
- Provide timely reporting to the LIS/EMR for clinician access.
- Provide timely oversight by the laboratory.
- Consolidate laboratory reporting.
- Automate charge capture.

Why not implement an automated electronic interface for charting of these glucose results? Some may say it is too expensive, but when you examine the cost to implement this type of interface, it pays for itself in the first year.

When looking at the cost for acquisition and implementation of an automated electronic interface, you will need to account for the following components. These components should reflect a one-time initial cost. If any have been bundled into a cost-per-strip price, the glucose vendors would need to provide a detailed accounting to reflect initial acquisition costs.

- Point-of-care system hardware acquisition and installation
- Point-of-care system acquisition
- Point-of-care system result interface
- Point-of-care system ADT interface
- Point-of-care system vendor implementation resource time
- Point-of-care system customer implementation resource time
- LIS/EMR result interface acquisition and installation
- LIS/EMR ADT interface acquisition and installation
- LIS/EMR interface testing resource time

Once values have been assigned to each of these items, the total can be subtracted from

the annual cost of manual charting to determine the ROI from eliminating manual charting. To calculate each subsequent year's cost for the automated electronic interface, you will need to consider the following components:

- Point-of-care system hardware maintenance.
- Point-of-care system and interface software maintenance.
- LIS/EMR result and ADT interface software maintenance.

With an automated electronic interface, the errors of omission and result transcription have been eliminated. However, some results will fail to post to the LIS/EMR because of an invalid patient ID entered on the device at test time, and time will now be required of the laboratory point-of-care team for the oversight and correction of these results. The amount of time spent performing just this task can also be determined by using the same formula used to calculate the time spent manually charting simply by substituting the number of results each day that require patient ID correction, the average amount of time required to address each result, and the average hourly rate for the individuals addressing this activity.

For example, using an average of a 5% daily error rate, 15 minutes per result, and \$40 per hour, the annual cost to address invalid patient ID errors on these results would be \$50,000. However, with bar-code scanning and positive patient ID on the meters at the time of testing, hopefully the number of results falling into this category can be decreased from the 5% used in this example.

Stepping away from glucose for the moment and looking at manual results that do not have an electronic output from the "device." It is fair to say the majority of hospitals perform manual testing and these results are being charted in one of the following ways:

1. Manually entered onto the patient's paper chart.
2. Manually entered into the EMR's nursing notes or flow sheet.
3. Manually written on log sheets, which the laboratory in turn enters into the LIS/EMR.
4. Manually written on log sheets and not entered into the LIS/EMR.



Referring back to the same article referenced in the beginning that addresses charting in the first two options above. There is an inherent delay with the third option and omission from the patient's record in the fourth option. Wouldn't it make more sense to provide a mechanism for the charting of manual results that provides the following benefits?

- Discrete results that can be queried
- Charting of both patient and liquid/aqueous quality control results
- Operator certification management
- Consistent reporting to the LIS/EMR
- Automated opportunities for charge/cost capture
- Timely laboratory oversight
- Complete customization to satisfy both laboratory and nursing

There are mechanisms provided by a couple of bedside glucose testing manufacturers and web applications provided by point-of-care middleware vendors.

As you take a moment to consider the inefficiencies of manual charting, it is important to take a step back and look at the bigger picture of your organization's current processes. You can do that by following these steps:

1. Gain a full understanding of what manual charting is being done. Think beyond glucose. What other point-of-care devices are being used that can be interfaced? What other departments (i.e., respiratory) are using bedside or near-patient testing? How are you electronically charting (or are you) results from manual tests? How accessible are your point-of-care results data?
2. Evaluate the costs associated with manual charting. When looking at this, consider both the actual financial costs and the value associated with having more timely and accurate access to the results, having the ability to run more timely and accurate reports, and providing patients with more complete electronic records.

3. Evaluate the costs associated with adding a solution that allows you to electronically interface the point-of-care testing results to the LIS/EMR, taking into account both the one-time costs and the annual maintenance costs.
4. Determine what your organization's ROI would be. How long would it take to "break even"? How much would you continue to save in subsequent years? By using the formulas discussed in this article, you can evaluate each of your point-of-care devices and see how the cost savings could add up over time in terms of actual staff productivity as well as the point-of-care data management efficiency, accuracy, and accessibility.

Maintaining accurate electronic records is an important way for you to provide quality care to your patients in a timely manner. By eliminating as much manual labor as possible through automatic electronic interfaces and having the ability to have manually entered results interfaced to the LIS/EMR, you can not only save time and money, but also ultimately enhance the patient experience. This article focused on the single aspect of manual charting and the ROI with automated electronic interfaces. However, there are also other significant benefits from centralized operator management and oversight that can contribute to the ROI and support accreditation compliance for operators performing manual charting.

.....

¹Enhancing Patient Safety Through Electronic Medical Record (EMR) Documentation of Vital Signs (Pauline Gearing, BSN, RN, CCRC; Christine Olney, MS, RN, doctoral candidate; Kim Davis, MS, ARNP, Diego Lozano, MD, Laura B Smith, MSN, RN, CCRN; and Bruce Friedman, D Eng).

Point of care & Volume 12, Number 2, June 2013
Reprints: Becky Clarke, BS, RN, TELCOR Inc, 7101 A St, Lincoln, NE 68510. Email: marketing@telcor.com.
The author declares no conflict of interest.
This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives 3.0 License, where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially. Copyright©2013 by Lippincott Williams & Wilkins.

